


# Long-term satellite data records of HCHO, H<sub>2</sub>O and CHOCHO


Introduction to OVOC/water vapor datasets for MEaSUREs

Christopher Chan Miller, Gonzalo Gonzalez Abad, Caroline Nowlan, Helen Wang, Kang Sun, Ewan O'Sullivan Xiong Liu, Kelly Chance

29th August 2019

# MEaSUREs will build on existing SAO algorithms to retrieve targets using observations from 6 instruments





#### Base code for all algorithms will be shared with TEMPO


Product will include algorithm traceability chains and error characterization borrowing from past projects e.g. QA4ECV (Boersma et al., 2018)

Instrument Retrievals done in two phases:

- (1) OMI, GOME-2, OMPS (overlapping validation data)
- (2) GOME, SCIAMACHY (validation via correlation with products from (1)




### Reliable long-term HCHO/CHOCHO observations will improve emissions attribution through temporal correlation



Zhu et al. (2017)




# Reliable long-term HCHO/CHOCHO observations will improve emissions attribution through temporal correlation



Zhu et al. (2017)



#### Water will be used as a fiducial reference for instrument harmonization




Wang et al. (2016)


Good-quality long term records exist for the entire MEaSUREs period enabling inter-sensor validation



### HCHO and CHOCHO will be indirectly validated using CTMs combined with aircraft observations



### HCHO and CHOCHO will be indirectly validated using CTMs combined with aircraft observations




Key Datasets


Algorithm

Level 1 Radiance
Radiance Reference "I<sub>0</sub>"


Spectrum Fit
Algorithm


Derive Slant Column via direct spectral fit using TEMPO codebase






















#### Testing the new stand-alone AMF Simulator



Next: Correct cloud products to account for inconsistent algorithm assumptions

Observations:
Radiance
O2 Slant Column




State Elements:
Surface Refl. (MODIS,Glint,Snow)
Aerosol

Lambertian Cloud Fraction/Pres.

#### Improved radiance references using cloudy targets




The typical radiance reference sector lies outside the TEMPO FOR




Using Clouds can shield boundary layer concentrations




#### Improved radiance references using cloudy targets





Radiances averaged for fully cloudy pixels between 700-800hPa

Radiances become stable after ~1 week averaging with high estimated precision (1x10-5)






#### **Project Timeline**


Initial release will be on the SAO Data Repo:

2021: OMI,GOME-2,OMPS

**2023: GOME, SCIAMACHY** 

Final Release Hosted on GES-DISC





Release will include daily Level 3 Products with new physics-based oversampling with archived grid weights

#### Acknowledgments:

NASA support through ACMAP Aura, TASNPP, and MEaSURES OMI/Aura and TEMPO Science Teams

